Entrez-PubMed

Page 1 of 2







Taxonomy OTABLE Structure PopSet Name Geromo Search PubMed Go Clear for Limits Preview/Index History Clipboard About Entrez Save Text Order Add to Clipboard Display Abstract

Entrez PubMed Overview Help (FAO)

☐ 1: Pancreas 1989;4(3):282-8

Related Articles, Books

Tulonal NEW Hew/Noteworthy

PubMed Services Journal Browser MeSH Browser Single Citation Malcher **Batch Citation** Matcher Clinical Queries

Cubby

Related Resources **Order Documents** Graleful Med Consumer Health Clinical Alerts Clinical trials.gov

Privacy Policy

Peptide YY: intrapancreatic localization and effects on insulin and glucagon secretion in the mouse.

Bottcher G, Ahren B, Lundquist I, Sundler F.

Department of Medical Cell Research, Lund University, Sweden.

We studied the intrapancreatic localization of peptide YY (PYY) and the effects of PYY on insulin and glucagon secretion in the mouse. Immunofluorescence staining of mouse pancreatic tissue showed that PYY occurred within islet cells. These cells were located preferentially at the periphery of the islets. Sequential and simultaneous double immunostaining revealed that most PYY cells also displayed glucagon immunoreactivity; some PYY cells contained immunoreactive pancreatic polypeptide (PP). At the electromicroscopic level, PYY immunoreactivity was demonstrated within the secretory granules of both glucagon cells and of a small granular cell type, which showed structural similarities to PP cells. In in vivo experiments, PYY at dose levels between 0.53 and 8.5 nmol/kg had no influence on basal plasma levels of insulin, glucagon, or glucose. In contrast, insulin secretion stimulated by glucose or the cholinergic agonist carbachol was inhibited by PYY (by 33 and 26%, respectively, at 4.25 nmol/kg). Similarly, carbachol-induced glucagon secretion was inhibited by PYY (by 47% at 4.25 nmol/kg). We conclude that PYY occurs in islet cells of the mouse pancreas, most of which are glucagon cells, and that PYY inhibits stimulated insulin and glucagon secretion in vivo in the mouse.

PMID: 2660131 [PubMed - indexed for MEDLINE]

Order Add to Clipboard Display Abstract Save Text


> Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Freedom of Information Act | Disclaimer

02.084

Clinical Alerts

Privacy Policy

ClinicalTrials.gov



PMID: 1513807 [PubMed - indexed for MEDLINE]

is exerted through a direct islet action.

Display Abstract Order Add to Clipboard Save Text

> Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Freedom of Information Act | Disclaimer

biphasic insulin response to arginine (10 mM). PYY also markedly reduced

conclude that in the perfused rat pancreas, PYY inhibits insulin secretion and induces vasoconstriction without a causal relationship. In addition, our results on isolated islets suggest that the inhibitory action of PYY on insulin secretion

the pancreatic vascular flow rate; this effect was observed at all three concentrations tested in a dose-dependent manner. In isolated islets, glucose (15 mM)-stimulated insulin secretion was inhibited by PYY at 10(-7) M. We